

V-BEAM LASER

BLOOD VESSEL LASER

SINGLE TREATMENTS


<i>Area(s)</i>	<i>Signature</i>	<i>Elite</i>
Face	\$450	\$550
Cheeks	\$330	\$430
Nose	\$300	\$400
Chin	\$300	\$400
Cheeks & Nose	\$350	\$450
Cheeks & Chin	\$350	\$450
Cheeks, Nose, & Chin	\$400	\$500
Face & Neck	\$775	\$875
Neck & Chest	\$775	\$875
Spot Treatment	\$300	\$400

PACKAGE OF 4

<i>Area(s)</i>	<i>Signature</i>	<i>Elite</i>
Face	\$1,530	\$1,870
Cheeks	\$1,122	\$1,462
Nose	\$1,020	\$1,360
Chin	\$1,020	\$1,360
Cheeks & Nose	\$1,190	\$1,530
Cheeks & Chin	\$1,190	\$1,530
Cheeks, Nose, & Chin	\$1,360	\$1,700
Face & Neck	\$2,635	\$2,975
Neck & Chest	\$2,635	\$2,975
Spot Treatment	\$1,020	\$1,360

TARGETS

Telangiectasias & Blood Vessels
Rosacea
Cherry Angiomas
Vascular Birth Marks
Bruises



*If you require a prescriptions pain management please contact the clinic 48 hours prior to treatment.
All prices subject to change without notice & may vary depending on individual needs*

PHONE
Call or text us
at 385.273.3376

SMART *Skin*
DERMATOLOGY

LOCATION
3200 W Clubhouse Dr. #100
Lehi, UT 84043

V-BEAM LASER

BLOOD VESSEL LASER

ABOUT TREATMENT

The Pulsed Dye Laser, or PDL is the gold standard for treatment of facial redness & superficial blood vessels. The laser specially targets hemoglobin, the major component of red blood cells. This in turn causes the blood vessel to collapse and fade naturally with time.

BEFORE TREATMENT

- **Sun Avoidance:** leading up to your treatment is important. We cannot treat patients who have an active sun burn or have too much recent sun exposure.
- **Numbing cream is not necessary:** The pain associated with a PDL treatment is fairly minimal, much like a light rubber band snap. The majority of patients will not require numbing cream, but a prescription of numbing may be called in at patient's request. Please note that numbing cream can sometimes flush the skin making for a more aggressive treatment.

AFTER TREATMENT

- Immediately following treatment you will likely see redness and swelling. and bruising. This will typically be the worst for the first 3-5 hours and will begin to fade.
- Possible bruising. Depending on the settings used and condition being treated, bruising may occur and is very normal. The bruising is usually accompanied by some increased swelling. Bruising can last 2-3 weeks.
- You may have been provided with a sample of triamcinolone (steroid) cream. If so, apply twice a day for 3-5 days to help decrease inflammation. Do not apply for a period of longer than one week.
- Apply gentle moisturizer twice a day as long as the bruising/red discoloration (purpura) is present and once per day if there is no discoloration or scabbing. It is important to keep the area moist until healed.
- Icing is recommended following the procedure. Icing over the first 24 hours will help decrease inflammation. Icing should be done in 15 minutes on, 15 minutes off intervals.
- Any discomfort you may have (usually not lasting more than a few hours, if any) can be relieved with acetaminophen (Tylenol).
- Avoid taking a bath during the first 48 hours as the intense heat may cause discomfort. Comfortably warm showers are recommended. Gently pat the area dry. Do not rub with a towel or washcloth.
- Do not rub, scratch, or pick at the treated area if bruising/red discoloration (purpura) is present.
- Avoid exposure to the sun. If sun exposure is expected, apply an SPF 30 or higher sunblock.

POSSIBLE SIDE EFFECTS

- **Common:** Redness, Swelling, Bruising
- **Rare:** Scarring, Hyperpigmentation, Blisters

Suggested Products: Tizo3 spf, EltaMD UV Clear or UV Daily spf. Gentle moisturizer such as Elta MD PM Therapy. Gentle cleanser such as Elta MD Foaming Facial Cleanser.

More Info:



Resources:

