
# SCLEROTHERAPY

LEG VEIN TREATMENT

## NUMBER OF TREATMENTS



SCLEROTHERAPY PRICING		
Visit Type	Signature Price	Elite Price
Standard Visit	\$250	\$350
Extended Visit	\$450	\$575

If you require a prescriptions pain management please contact the clinic 48 hours prior to treatment.

All prices subject to change without notice & may vary depending on individual needs



# SCLEROTHERAPY

LEG VEIN TREATMENT

#### LEADING UP TO TREATMENT

- Compression Stockings. These will need to be purchased either at a medical supply store or Online. 20/30 mm Hg strength is recommended. Stockings will need to be long enough to cover all treated
- veins. Knee high, thigh high can be used, but typically our patients find full length stockings the most comfortable and easy to use.
- Wear loose fitting shorts to your visit. These are typically easier and more comfortable than a gown.
- Avoid wearing lotion on legs on day of procedure.

#### WHAT TO EXPECT

- Pain. You may experience mild stinging or aching during and post injection.
- Itching. The injected solution acts as an irritant to the blood vessel. Some individuals may experience some mild itching as a result.
- Darkening of vessels. This is part of the normal treatment course. As the blood begins to clot and the vessel collapses, it will take on a darkened appearance before lightening.
- "Trapped blood". This may feel like a sore lump and occurs often in large varicose veins after treatment. We typically recommend draining the site with a small puncture which is done in the office.
- "Blushing". This is the appearance of new tiny spider veins around an injection site. This occurs in roughly 25% of treated patients and will resolve naturally without additional treatments.

### AFTER TREATMENT

- Wear your compression stockings for 14 days following your treatment. They can be taken off to shower. We suggest wearing them while sleeping but if this is too difficult it is okay to have them off.
- Icing after your procedure in 15 minute on, 15 minute off intervals for the first 24 hours can help with
- swelling or pain.
- Avoid Motrin or Advil. Take over the counter Tylenol as needed for discomfort.
- Take over the counter Benadryl as needed for itching.
- Avoid any strenuous exercise or weight lifting for 7 days following procedure.
- No airplane rides for at least 14 days after treatment.
- Do not expose treated area to the sun or tanning bed.

### POSSIBLE SIDE EFFECTS

- Common: Redness, Swelling, bruising, pain with injection, matted blood vessels, itching
- Rare: blood clot, allergic reaction, sores

Suggested Products: EltaMD AOX Mist sunscreen





Resources:



