


PRF FOR HAIR LOSS

COLLAGEN STIMULATING HAIR GROWTH TREATMENT



TARGETS

Hair Loss
Follicle stimulation



S	Signature	E	Elite
	ELIZABETH HARDING NP-C		

PRF 6-26

SINGLE TREATMENT		
E	1 vial	\$750
F	2 vials	\$1,250

S	Signature	E	Elite
	ELIZABETH HARDING NP-C		

PRF 6-26

PACKAGE OF 3 TREATMENTS		
E	1 vial each	\$1,900
F	2 vials each	\$3,150

If you require prescription pain management please contact the clinic 48 hours prior to treatment.

All prices subject to change without notice & may vary depending on individual needs.

Number of ml will vary patient to patient dependant on blood drawn, spun, and injected during one visit.

PHONE
Call or text us
at 385.273.3376

SMART Skin
DERMATOLOGY

LOCATION
3200 W Clubhouse Dr. #100
Lehi, UT 84048

PRF FOR HAIR LOSS

COLLAGEN STIMULATING HAIR GROWTH TREATMENT

ABOUT TREATMENT

Platelet-Rich Fibrin (PRF) is an innovative treatment that uses your body's own regenerative abilities to address hair loss. With a system like Selphyl® PRFM, a small blood sample is taken, much like a standard blood test. This sample is then carefully processed to create a concentrated solution of your platelets and fibrin. These components are packed with growth factors crucial for tissue repair and regeneration.

This natural, golden-yellow solution is then skillfully injected into areas of the scalp where hair thinning or loss is present. The fibrin acts as a natural scaffold, holding the platelets in place and allowing for a slow, sustained release of these beneficial growth factors. This prolonged release works to stimulate dormant hair follicles, promote new hair growth, and improve the overall health and thickness of existing hair.

BEFORE TREATMENT

- Drink 1-2 electrolyte drinks 24 hours before your treatment. Your hydration level will dictate how much product is obtained. A well hydrated patient will get 1 ml more PRF on average.
- Arnica Montana: If you have a history of bruising with injections. This is a supplement that can help prevent or lessen bruising. should be started 1-2 day(s) prior to treatment for best effect.
- Aspirin / Ibuprofen avoidance: If possible, skip Aspirin 2 weeks prior to your treatment. Avoid Ibuprofen or Aleve 3 days prior to procedure.

AFTER TREATMENT

- Ice on the affected area post-procedure in 15 minutes on, 15 minutes off intervals for the first 24 hours, can help with swelling or pain.
- Avoid any rubbing or massaging of injection sites unless expressly instructed by your physician.
- If doing package of PRF, return in 4-6 weeks for your next treatment.

POSSIBLE SIDE EFFECTS

- **Common:** Redness, swelling/bruising, pain at injection site
- **Rare:** Formation of granulomas, and skin necrosis

More Info:



Resources:

