

WART PEEL

PATIENT INFORMATION & INSTRUCTIONS

PRESCRIBERS

Please fax prescriptions to (319) 354 - 6050 or E-Prescribe following these instructions: www.wartpeel.com/eprescribing

WartPEEL treats the following conditions (check all that apply):

☐ Warts

☐ Genital Warts

☐ Seborrheic Keratosis

☐ Molluscum



1900 James Street, Ste. 10
Coralville, IA 52241
Ph: (319) 354 - 6006
Hours:
Mon-Fri: 9 AM-5:30 PM CST

PATIENTS

You have been prescribed a compounded medication from Medicap Pharmacy. They will be in touch with you soon in regards to your prescription.

HOW IT WORKS

1. Patient will receive a text or email once the pharmacy processes the prescription.
2. A link will be provided that will direct the patient to the shopping cart
3. Patient answers necessary medical questions, watches educational video, and selects mail-out option.
4. Patient then completes the payment process. Major credit cards and FSA/HSA payment methods are accepted!



Patient instructions can be found at wartpeel.com/for-patients. Scan the QR Code to quickly find written and video instructions for the condition you are using WartPEEL to treat.

OTHER RESOURCES

We work hard to publish only the best patient education and comprehensive resources. We want to be your trusted source of all skin information. You will find additional educational information by scanning the following QR codes.

*Our Monthly
Newsletter*



*Patient
Resources*



PHONE
Call or text us
at 385.273.3376

SMART *Skin*
DERMATOLOGY

LOCATION
3200 W Clubhouse Dr. #100
Lehi, UT 84048

WART PEEL

PATIENT INFORMATION & INSTRUCTIONS

INSTRUCTIONS

STEP 1: Apply once daily at bedtime. Remove tip cap and apply a small amount of WartPEEL directly onto the plastic applicator (or a toothpick). Use forward and backward pressure on the plunger to control the amount of medication that comes out of the tip of the syringe. Replace tip cap to prevent the medication from drying out.

STEP 2: WartPEEL is a glue not a cream, so make sure it fully dries. After applying a thin and even film on the wart or other lesions being treated, wait until it dries completely (10 to 20 minutes) and a color change from white to translucent white occurs. Test dryness with the blue applicator. WartPEEL should not stick to the applicator and is slightly soft when fully dry. The drying process can be sped up by using a hair dryer on the cool setting or a fan. Be careful not to get the medication onto the healthy skin as this may lead to irritation.

STEP 3: Apply a small piece of the 3M Blendederm tape that came with WartPEEL over the fully dry medication and leave on overnight. Tape is only recommended for use on the hands, feet, elbows, and knees unless otherwise directed by your health care professional.

STEP 4: Wash hands after applying the medication. Never apply to the mouth, eyes, or inside the nose.

STEP 5: In the morning, remove the tape and wash the area to remove excess medication. The area can be left uncovered during the day or a covering such as a band aid may be used.

STEP 6: Repeat steps 1-5, applying and removing the medication once daily for 1-3 weeks or as directed.

STORAGE

Store this medication at room temperature. Keep out of reach of children.

Expiration Date:

The medication is best used within four months of the date it was made. Do not use or keep outdated medication.

Side Effects

- Redness and irritation around the wart(s) are most commonly reported
- Irritation can be minimized with careful application to the wart alone
- Irritation can be minimized by allowing the medication to dry completely before applying tape
- Though uncommon, burning, crusting, allergic contact dermatitis, itchiness, scarring, rash, soreness, and ulceration can occur.
- Call your doctor or seek medical attention if you have any side effects that bother you or do not go away.

Contraindications

- Do not use if you are pregnant or breastfeeding
- May cause fetal harm when given to pregnant women
- Medications present in breast milk may lead to life-threatening or fatal side effects in nursing babies
- Do not use if you have a severe allergic reaction (anaphylaxis) to WartPEEL or its components including:
- Salicylic acid or derivatives such as aspirin
- 5-fluorouracil (5-FU)